

01. Mala of Medo dhatu is :

- | | |
|-------------|---------------|
| (1) Mutra | (2) Sveda |
| (3) Purisha | (4) Medo roga |

02. Takrarista is indicated in which disease ?

- | | |
|----------------|---------------|
| (1) Sandhivata | (2) Gridhrasi |
| (3) Urustambha | (4) Grahani |

03. The synonym of Rasakriya is :

- | | |
|------------|-------------|
| (1) Kalka | (2) Avaleha |
| (3) Churna | (4) Vati |

04. According to Charak Samhita, Sankhapushpi should be used as Medhya Rasayana in form of :

- | | |
|-------------|------------|
| (1) Svarasa | (2) Churna |
| (3) Kalka | (4) Kwatha |

05. Main Kalpana of Snehapana is :

- | | |
|-----------|---------------|
| (1) Vasti | (2) Yusha |
| (3) Audan | (4) Acchapeya |

06. Use of Kalyanaka Ghrita is described in Kashyapa Samhita in :

- | | |
|---------------------------|---------------------------|
| (1) Vedanadhayaya | (2) Lehadhayaya |
| (3) Dantajanmika adhayaya | (4) Visesa Kalpa adhayaya |

07. Vamana is the choice of treatment in which type of ajirna :
- | | |
|--------------------|-------------------|
| (1) Amajirna | (2) Vidagdhajirna |
| (3) Vistabdhajirna | (4) Rasasesajirna |
08. According to Charak Samhita, pure blood resembles the colour of :
- | | |
|--------------------|----------------------|
| (1) Gunja phala | (2) Bhallataka phala |
| (3) Udumbara phala | (4) Madhuka phala |
09. Which of the following drug is **not** Yogavahi ?
- | | |
|-------------|-------------|
| (1) Draksha | (2) Pippali |
| (3) Madhu | (4) Ghrita |
10. Gunja is enumerated in Gana called :
- | | |
|---------------|---------------|
| (1) Upavisha | (2) Visha |
| (3) Mahavisha | (4) Garavisha |
11. Which of the following hernias follows the path of the spermatic cord within the cremaster muscle ?
- | | |
|-----------------------|---------------------|
| (1) Femoral | (2) Direct inguinal |
| (3) Indirect inguinal | (4) Spigelian |
12. A 76-year old man is scheduled to undergo repair of an abdominal hernia that is easily reduced by pushing the abdominal contents back through the external ring. During repair, the surgeon sees that the hernial sac protrudes from the abdominal wall superior to the inguinal ligament and medial to the inferior epigastric vessels. Which of the following types of hernia does this patient have ?
- | | |
|--------------------------|------------------------------|
| (1) Diaphragmatic hernia | (2) Direct inguinal hernia |
| (3) Femoral hernia | (4) Indirect inguinal hernia |

13. Most common intra-abdominal tumor in children :
- (1) Neuroblastoma (2) Wilm's Tumour
(3) Lymphoma (4) Sarcoma
14. The best indication of ERCP is :
- (1) Obstructive jaundice (2) History of jaundice
(3) Gallstone pancreatitis (4) 10mm CBD stone seen on US
15. Regarding malignant tumour of appendix :
- (1) RT hemicolectomy is the treatment for all carcinoid tumour
(2) Carcinoid tumor of the appendix is the 2nd most common malignancy
(3) Appendix with carcinoid lowers 5y survival than non carcinoid appendix.
(4) Apendix goblet carcinoid< 1cm is safely treated with simple appendectomy
16. Which of the following is **not** a cancer sign ?
- (1) gray turner sign (2) krockenbuerge
(3) virchow sign (4) sister marry joseph
17. All of the following form radiolucent stones except :
- (1) Xanthine (2) Cysteine
(3) Allopurinol (4) Orotic acid

18. The direction of flow of venous blood in conditions of valve incompetence affecting perforating veins of lower limb is ?
- (1) along gravity (2) superficial to deep
(3) along osmotic gradient (4) deep to superficial
19. Structures that do not come during surgery for vasectomy :
- (1) Testicular artery
(2) Ilioinguinal nerve
(3) Pampiniform plexus
(4) Genitofemoral branch of femoral nerve
20. Which is one of the lekhnaya disease ?
- (1) Vidradhi (2) Ashmari
(3) Granthi (4) Balmika
21. Seevan karma is contraindicated in :
- (1) medaj vrana (2) bhinna vrana
(3) marutvahin vrana (4) sadhyo vrana
22. Gada Bandhan is not indicated in :
- (1) kaksha (2) vankshan
(3) shakha (4) head
23. Kehr's sign is positive in case of :
- (1) ruptured spleen (2) acute cholecystitis
(3) nephritis (4) acute pancreatitis

24. E R C P is the diagnostic & therapeutic for :

- | | |
|--------------------|------------------------|
| (1) Pancreatitis | (2) Cholecystitis |
| (3) Cholelithiasis | (4) Choledocolithiasis |

25. Murphy's sign positive is found in case of :

- | | |
|------------------------|-------------------------|
| (1) acute appendicitis | (2) acute cholecystitis |
| (3) acute pancreatitis | (4) acute peritonitis |

26. Whipple's operation is performed in case of :

- | | |
|--------------------|-------------------------|
| (1) Ca Stomach | (2) ca gall bladder |
| (3) nephroblastoma | (4) ca head of pancreas |

27. Billroth I operation is indicated for :

- | | |
|--------------------|-------------------------|
| (1) pyloric growth | (2) duodenal ulcer |
| (3) gastric ulcer | (4) ca head of pancreas |

28. Bassini's repair of hernia is appropriate for :

- | | |
|------------------------------|----------------------------|
| (1) congenital hernia | (2) inguinal direct hernia |
| (3) inguinal indirect hernia | (4) femoral hernia |

29. Gaillard Thoms's method of breast operation is indicated in a case of :

- (1) young women have begun breast tumour
- (2) ca breast
- (3) duct papilloma
- (4) large breast abscess

30. Anderson-Hynens operation is indicated for the a case of :
- (1) nonfunctioning kidney (2) pyonephrosis
(3) cong hydronephrosis (4) horse shoe kidney
31. In portal hypertension the sites of portosystemicanastamosis include :
- (1) Lower end of esophagus,
(2) Around umbilicus,
(3) Lower third of rectum and anal canal,
(4) All of the above
32. In Budd chiari syndrome the occlusion is at the :
- (1) IVC, (2) Renal vein,
(3) Hepatic vein, (4) Splenic vein
33. Line of surgical division of the lobes of the liver is :
- (1) Falciform ligament to the diaphragm,
(2) Gall bladder bed to IVC,
(3) Gall bladder bed on the Lt. crus of diaphragm,
(4) One inch to the left of falciform ligament to the IVC
34. Ectopic testis is found in all locations except :
- (1) Lumbar, (2) Perineal,
(3) Intra abdominal, (4) Inguinal

35. Treatment of Zollinger Ellison syndrome :

- (1) Total gastrectomy with removal of tumour,
- (2) Partial gastrectomy,
- (3) Excision of tumour alone
- (4) H₂ receptor antagonist

36. Earliest symptoms of carcinoma Rectum :

- (1) Pain
- (2) Alteration of bowel habits
- (3) Bleeding PR
- (4) Tenesmus

37. Staghorn calculus is made out of :

- (1) Oxalate
- (2) Phosphate,
- (3) Uric acid,
- (4) Cystine

38. Honey comb liver is seen in :

- (1) Micronodular cirrhosis,
- (2) Dubin Johnson's syndrome,
- (3) Actinomycosis,
- (4) Hydatidosis

39. In radical mastectomy the structures preserved are all except :

- (1) Axillary vein,
- (2) Cephalic vein,
- (3) Nerve to serratus anterior,
- (4) Pectoralis minor

40. Indications for surgery in Benign prostatic hypertrophy are all except :

- (1) Prostatism,
- (2) Chronic retention
- (3) Hemorrhage,
- (4) Enlarged prostate